

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/445, 9/00	A1	(11) International Publication Number: WO 96/11687 (43) International Publication Date: 25 April 1996 (25.04.96)
(21) International Application Number: PCT/EP95/03966 (22) International Filing Date: 6 October 1995 (06.10.95) (30) Priority Data: 94202986.9 14 October 1994 (14.10.94) EP (34) Countries for which the regional or international application was filed: AT et al. (71) Applicant (for all designated States except US): JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): FRANÇOIS, Marc, Karel, Jozef [BE/BE]; Foxemastraat 64, B-2920 Kalmthout (BE). AGEMANS, Christine, Frieda, Augusta [BE/BE]; Oudstrijdersstraat 24, B-2520 Oelegem (BE). (74) Agent: QUAGHEBEUR, Luc; Janssen Pharmaceutica N.V., Patent Dept., Turnhoutseweg 30, B-2340 Beerse (BE).		(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, HU, IS, JP, KE, KR, LT, LV, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TT, UA, UG, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: SABELUZOLE ORAL SUSPENSIONS (57) Abstract An aqueous suspension for oral administration comprising sabeluzole and a pharmaceutically acceptable carrier, having a pH in the range from 8 to 10; processes for preparing the same.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

SABELUZOLE ORAL SUSPENSIONS

5

The invention relates to physicochemically stable sabeluzole formulations having a satisfactory taste and aftertaste.

10

In US-4,861,785 there are described compounds having antihypoxic and antianoxic properties useful in indications such as shock, cardiac arrest and severe blood loss. Among these compounds features 4-(2-benzothiazolylmethylamino- α -(4-fluorophenoxy)methyl)-1-piperidineethanol, which is generically known as sabeluzole.

15

Subsequent investigations have shown the potential of sabeluzole in the treatment of patients suffering from chronic neuro-degenerative diseases such as dementia of the Alzheimer type (DAT) or Alzheimer's disease, amyotrophic lateral sclerosis (ALS), dementia associated with Parkinson's disease and other central nervous system diseases which are characterized by progressive dementia.

20

Administration of an oral dosage form is the preferred route of administration for many pharmaceuticals because it provides for easy, low-cost administration. However, patient compliance can be a problem when the patient is requested to swallow a solid formulation such as a tablet or a capsule. Therefore, the development of a liquid oral formulation is often desirable.

25

The development of a liquid oral formulation of sabeluzole is hampered by the unpleasant bitter taste of the compound. The bitter flavour of sabeluzole is inevitably tasted during drinking or immediately after swallowing a solution comprising the compound. A clear improvement in taste is observed using a suspension of sabeluzole instead of a solution of the compound. In particular the bitter aftertaste is significantly reduced. The development of a useful suspension of sabeluzole, however, is hampered by chemical stability problems of the compound. In particular, aqueous suspensions of sabeluzole at pH 7 suffer from degradation of sabeluzole and have an unacceptable shelf-life. A stable sabeluzole suspension was prepared by maintaining the pH in a strict range from about 8 to 10. These suspensions were further specifically adapted so as to allow them to be diluted with cold beverages such as fruit juice and also hot beverages such as tea, cocoa and coffee. These features are considered essential properties in a medicament which has to be administered to patients whose compliance with therapy is a major concern.

40

The present invention is concerned with stable aqueous sabeluzole suspensions having a pH in the range from 8 to 10. In particular, the invention relates to aqueous suspensions for oral administration comprising sabeluzole and a pharmaceutically acceptable carrier, having a pH in the range from 8 to 10.

5

The term "stable" as used herein relates to compositions wherein the decrease in the sabeluzole content is less than 10%, preferably less than 5% and most preferably less than 2%, after storage at room temperature for up to 3 months.

10 The term sabeluzole as used hereinabove also comprises the solvates which sabeluzole is able to form and said solvates are meant to be included within the scope of the present invention. Examples of such solvates are, e.g. the hydrates, alcoholates and the like.

15 Sabeluzole has an asymmetric carbon atom and the absolute configuration of this asymmetric centre may be indicated by the stereochemical descriptors R and S. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, in particular the racemate.

20 Two polymorphs of sabeluzole base are known. The higher melting polymorph I (mp. 101.7°C, maximum 105.3°C, heat of fusion = 99.6 J/g) can be distinguished easily by its DSC characteristics from the lower melting polymorph II (mp. 88.9 °C, maximum 91.6°C, heat of fusion 84.2 J/g). X-ray diffraction analysis has confirmed the existence of the two polymorphs. Polymorph II is metastable and dissolves considerably more rapidly than polymorph I, especially at mildly acidic pH values. Polymorph II when
25 substantially crystallographically pure is stable for all practical purposes. Preferably, the suspensions include the stable polymorph I.

Hereinafter, the amounts of each of the ingredients in the compositions are expressed as percentages by weight based on the total volume of the formulation, unless otherwise
30 indicated.

In particular, the concentration of sabeluzole in the present suspensions may range from 0.01% to 5%, preferably from 0.05% to 1%, more preferably from 0.1% to 0.5% and in
35 particular is about 0.1%.

Chemical degradation of sabeluzole in suspension is prevented by raising the pH to slightly to moderately basic, that is to pH 8 to 10, in particular to pH of approximately 9. A maximum trade-off between two mutually contrary prerequisites, namely increasing chemical stability with increasing pH and decreasing organoleptic properties with

increasing pH, is reached at about pH 9. In view of the further prerequisite that the suspension should be dilutable with a variety of beverages, said pH ranges are created by using a buffer system. Buffer systems comprise mixtures of appropriate amounts of an acid such as phosphoric, hydrochloric or boric acid, and a base, in particular sodium carbonate, sodium bicarbonate, sodium hydroxide or disodium hydrogen phosphate. Said buffer systems should maintain the pH of the formulation in the range from 8 to 10, more preferably in the range from 8.5 to 9.5 and most preferably at about 9. Preferably, a carbonate buffer comprising sodium carbonate, sodium bicarbonate and/or sodium hydroxide or hydrochloric acid is used.

The oral suspension may further comprise various pharmaceutically acceptable ingredients such as suspending agents, wetting agents, stabilizing agents, preservatives, and the like. Suitable suspending agents are cellulose derivatives, e.g. dispersible cellulose (= a mixture of microcrystalline cellulose and carboxymethylcellulose sodium), hypromellose (= hydroxypropyl methylcellulose), and the like. Preferably, dispersible cellulose (Avicel RC591®) is used in an amount of 0.1 to 2%, more preferably in an amount of about 1.2%. The addition of Avicel RC591® results in thixotropic properties of the suspension, i.e. the suspension becomes temporarily liquid when shaken or stirred and returns to a gel on standing. The gel structure of the suspension upon standing precludes the suspended particles to precipitate. Hydrocolloids such as hypromellose stabilize the suspension by adsorption to the suspended particles. Hypromellose (hydroxypropyl methylcellulose) has proven to be particularly useful except when the suspension is to be diluted with a hot beverage because the agent tends to gel and flocculate. Fortunately, it has been found that the incorporation of hypromellose is entirely optional and that aqueous sabeluzole suspensions without hypromellose meet all the prerequisites set out hereinbefore and, in addition, can easily be diluted with hot beverages without the problem of flocculation of any of the adjuvants used in the formula. If used at all, an amount of 0.1% to 2.5% in particular about 0.25% (w/v), of hypromellose, is adequate.

Suitable wetting agents are, for example, polyoxyethylene derivatives of sorbitan esters, e.g. polysorbate 20 (Tween 20®), polysorbate 40 (= Tween 40®), polysorbate 60 (= Tween 60®), and the like. In particular, polysorbate 20 is used in an amount of 0.01 to 1%, preferably in an amount of approximately 0.025%.

Suitable preservatives which are stable at the alkaline conditions of the suspension are, e.g. propylene glycol, ethanol and the like. Preferably, propylene glycol is used in an amount of 5 to 30% (v/v), more preferably in an amount of about 20% (v/v).

In order to improve the palatability of the suspension sweeteners and/or flavouring substances may be added. Sweeteners have been found to improve the organoleptic properties of aqueous sabeluzole suspensions at pH 8 to 10 very markedly. Flavouring agents on the other hand appear to be entirely optional; while definitely influencing the taste of the suspension, they do not appear to improve the organoleptic properties of said suspensions. Suitable sweeteners comprise saccharin, sodium or calcium saccharin, aspartame, acesulfame potassium, sodium cyclamate, alitame, a dihydrochalcone sweetener, monellin, stevioside or sucralose (4,1',6'-trichloro-4,1',6'-trideoxy-galactosucrose), or a bulk sweetener such as sorbitol, mannitol, fructose, sucrose, maltose, isomalt, glucose, hydrogenated glucose syrup, xylitol, caramel or honey. In particular, polyalcohols are used as sweetening agents. The latter show the additional advantage that they increase the viscosity of the suspension and enhance the antimicrobial efficacy of, e.g. propylene glycol. Preferably, sorbitol (in a 70% w/v solution) is used in an amount of 5 to 30% (v/v), more preferably in an amount of about 20% (v/v).

Suitable flavours which may optionally be added are Chocolate flavour, Herb flavour, Caramel Chocolate flavour, Mint Cool flavour, Fantasy flavour and fruit flavours such as cherry, raspberry, black currant or strawberry flavour, and the like. Each flavour may be present in the final composition in a concentration ranging up to 1%. Combinations of flavours may advantageously be used. Obviously, the flavours used preferably do not undergo any change or loss of taste and colour under the alkaline conditions of the formulation.

The subject suspensions may be presented in art-known containers such as bottles, spray devices, sachets, and the like. Optionally, the suspensions are manufactured in unit-dose containers, e.g. unit-dose sachets or unit-dose bottles.

In general it is contemplated that an effective daily amount would be from 1 to 40 mg, preferably from 10 to 20 mg of active ingredient. It is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore guidelines only and are not intended to limit the scope or use of the invention to any extent.

In particular, the present invention relates to suspensions comprising:

- (a) 0.01 to 5% (w/v) sabeluzole;
- (b) 0.2 to 4% (w/v) suspending agents;
- (c) up to 30% (w/v) sweeteners;

-5-

- (d) up to 30% (v/v) preservatives;
- (e) 0.01 to 1% (w/v) wetting agent;
- (f) buffer substances to fix the pH in the range from 8 to 10; and
- (g) water q.s. ad 100%.

5

More particularly, the present invention is concerned with suspensions comprising:

- (a) 0.05 to 1% (w/v) sabeluzole;
- (b) 0.8 to 1.5% (w/v) dispersible cellulose and optionally 0.1 to 0.5% (w/v) hypromellose;
- 10 (c) 10 to 30% (v/v) sorbitol solution (70% (w/v) in water);
- (d) 10 to 30% (v/v) propylene glycol;
- (e) 0.01 to 1% (w/v) polysorbate 20;
- (f) buffer substances to fix the pH in the range from 8.5 to 9.5; and
- (g) water q.s. ad 100%.

15

Preferably, the invention relates to a suspension comprising approximately:

- (a) 0.1% (w/v) sabeluzole polymorph I;
- (b) 1.2% (w/v) dispersible cellulose and optionally 0.25% (w/v) hypromellose;
- (c) 20% (v/v) sorbitol 70% (w/w) solution;
- 20 (d) 20% (v/v) propylene glycol;
- (e) 0.025% (w/v) polysorbate 20;
- (f) sodium carbonate and hydrochloric acid to fix the pH at about 9;
- (g) water q.s. ad 100%

- 25 In a particular aspect of the invention the above suspensions may include one or more flavouring substances.

- Further, the present invention relates to the preparation of the described suspensions. The preparation involves the intimate mixing of the active ingredient with the carrier
- 30 ingredients. In particular, the preparation involves the following steps: (a) the suspending agents, wetting agents, sweeteners and preservatives are mixed with an amount of water; (b) sabeluzole is mixed with phase (a); and (c) the pH is fixed in the range from 8 to 10.

- 35 Optionally, the above procedure is conducted under an inert atmosphere, e.g. nitrogen or oxygen-free argon. It is advantageous to use a micronized form of sabeluzole, in particular material having an average particle size of less than 100 microns, preferably less than 75 microns, and in particular having a mean particle size of not more than 15

microns. Micronized forms can be prepared by micronization techniques known in the art, e.g. by milling in appropriate mills and sieving through appropriate sieves.

5 The following examples are intended to illustrate the scope of the present invention in all its aspects.

Example 1:

F1

	<u>Ingredient</u>	<u>Quantity</u>
10	Sabeluzole polymorph I	2.5 mg
	Polysorbate 20	0.25 mg
	Dispersible cellulose	12 mg
	Propylene glycol	200 µl
	Hypromellose 2910	2.5 mg
15	Sorbitol 70% (w/w) solution	200 µl
	Sodium carbonate	5 mg
	Concentrated hydrochloric acid	q.s. ad pH=9
	Purified water	q.s. ad 1ml

20 Preparation:

- (1) 2.5 mg hypromellose 2910 was added upon stirring to an amount of purified water at 90-95°C;
- (2) phase (1) was cooled to ambient temperature;
- (3) 200 µl sorbitol 70% (w/w) solution, 0.25 mg polysorbate 20, 200 µl propylene
- 25 glycol and 12 mg dispersible cellulose were homogeneously dispersed in an amount of purified water;
- (4) phases (2) and (3) were mixed;
- (5) 2.5 mg sabeluzole polymorph I was added to phase (4);
- (6) 5 mg sodium carbonate was dissolved in an amount of purified water;
- 30 (7) phases (5) and (6) were mixed upon stirring;
- (8) the pH of phase (7) was adjusted to about 9 with concentrated hydrochloric acid; and
- (9) phase (8) was diluted to the desired end volume.

In a similar way there were prepared:

-7-

F2:

	<u>Ingredient</u>	<u>Quantity</u>
	Sabeluzole polymorph I microfine	5 mg
	Polysorbate 20	0.25 mg
5	Dispersible cellulose	12 mg
	Propylene glycol	200 µl
	Hypromellose 2910	2.5 mg
	Sorbitol 70% (w/w) solution	150 µl
	Sodium carbonate	2.5 mg
10	Herb flavour	0.53 mg
	Concentrated hydrochloric acid	q.s. ad pH=9
	Sodium hydroxide	q.s. ad pH=9
	Purified water	q.s. ad 1ml

15 F3:

	<u>Ingredient</u>	<u>Quantity</u>
	Sabeluzole polymorph I microfine	5 mg
	Polysorbate 20	0.25 mg
	Dispersible cellulose	12 mg
20	Propylene glycol	200 µl
	Hypromellose 2910	2.5 mg
	Sorbitol 70% (w/w) solution	150 µl
	Sodium carbonate	5 mg
	Herb flavour	0.53 mg
25	Concentrated hydrochloric acid	q.s. ad pH=9
	Sodium hydroxide	q.s. ad pH=9
	Purified water	q.s. ad 1ml

F4:

	<u>Ingredient</u>	<u>Quantity</u>
30	Sabeluzole polymorph I microfine	5 mg
	Polysorbate 20	0.25 mg
	Dispersible cellulose	12 mg
	Propylene glycol	200 µl
35	Hypromellose 2910	2.5 mg
	Sorbitol 70% (w/w) solution	150 µl
	Sodium carbonate	3.18 mg
	Sodium bicarbonate	5.88 mg

-8-

	Herb flavour	0.53 mg
	Concentrated hydrochloric acid	q.s. ad pH=9.5
	Sodium hydroxide	q.s. ad pH=9.5
5	Purified water	q.s. ad 1ml

F5:

	<u>Ingredient</u>	<u>Quantity</u>
	Sabeluzole polymorph I	1 mg
	Polysorbate 20	0.25 mg
10	Dispersible cellulose	12 mg
	Propylene glycol	200 µl
	Hypromellose 2910	2.5 mg
	Sorbitol 70% (w/w) solution	200 µl
	Sodium carbonate	5 mg
15	Concentrated hydrochloric acid	q.s. ad pH=9
	Purified water	q.s. ad 1ml

F6 :

	<u>Ingredient</u>	<u>Quantity</u>
20	Sabeluzole polymorph I	2.5 mg
	Polysorbate 20	0.25 mg
	Dispersible cellulose	12 mg
	Propylene glycol	200 µl
	Sorbitol 70% (w/w) solution	200 µl
25	Sodium carbonate	5 mg
	Concentrated hydrochloric acid	q.s. ad pH=9
	Purified water	q.s. ad 1ml

F7:

	<u>Ingredient</u>	<u>Quantity</u>
30	Sabeluzole polymorph I microfine	5 mg
	Polysorbate 20	0.25 mg
	Dispersible cellulose	12 mg
	Propylene glycol	200 µl
35	Sorbitol 70% (w/w) solution	150 µl
	Sodium carbonate	2.5 mg
	Herb flavour	0.53 mg
	Concentrated hydrochloric acid	q.s. ad pH=9

-9-

Sodium hydroxide	q.s. ad pH=9
Purified water	q.s. ad 1ml

F8:

5	<u>Ingredient</u>	<u>Quantity</u>
	Sabeluzole polymorph I microfine	5 mg
	Polysorbate 20	0.25 mg
	Dispersible cellulose	12 mg
	Propylene glycol	200 µl
10	Sorbitol 70% (w/w) solution	150 µl
	Sodium carbonate	5 mg
	Herb flavour	0.53 mg
	Concentrated hydrochloric acid	q.s. ad pH=9
	Sodium hydroxide	q.s. ad pH=9
15	Purified water	q.s. ad 1ml

F9:

	<u>Ingredient</u>	<u>Quantity</u>
	Sabeluzole polymorph I microfine	5 mg
20	Polysorbate 20	0.25 mg
	Dispersible cellulose	12 mg
	Propylene glycol	200 µl
	Sorbitol 70% (w/w) solution	150 µl
	Sodium carbonate	3.18 mg
25	Sodium bicarbonate	5.88 mg
	Herb flavour	0.53 mg
	Concentrated hydrochloric acid	q.s. ad pH=9.5
	Sodium hydroxide	q.s. ad pH=9.5
30	Purified water	q.s. ad 1ml

F10:

	<u>Ingredient</u>	<u>Quantity</u>
	Sabeluzole polymorph I	1 mg
	Polysorbate 20	0.25 mg
35	Dispersible cellulose	12 mg
	Propylene glycol	200 µl
	Sorbitol 70% (w/w) solution	200 µl
	Sodium carbonate	5 mg

-10-

Concentrated hydrochloric acid
Purified water

q.s. ad pH=9
q.s. ad 1ml

Example 2

- 5 The suspensions as described hereinabove were stored for 3.5 months at room temperature. The concentration of sabeluzole had not significantly changed after storage. No degradation products were detected. Hence, the described suspensions are in compliance with the requirements of a stable formulation as set forth hereinabove.

Claims

1. An aqueous suspension for oral administration comprising sabeluzole and a pharmaceutically acceptable carrier, having a pH in the range from 8 to 10.
2. A suspension according to claim 1 wherein the active ingredient is sabeluzole polymorph I.
3. A suspension according to claim 1 wherein the pH range from 8 to 10 is created with a carbonate buffer comprising sodium carbonate, sodium bicarbonate and/or sodium hydroxide or hydrochloric acid.
4. A suspension according to claim 3 wherein the pH is approximately 9.
5. A suspension according to claim 1 further comprising a suspending agent.
6. A suspension according to claim 5 wherein the suspending agent is Avicel RC591® in an amount of 0.1 to 2% (w/v).
7. A suspension according to claim 1 further comprising a sweetener.
8. A suspension according to claim 7 wherein the sweetener is a sorbitol solution (70% (v/v)) in an amount of 5 to 30% (w/v).
9. A suspension according to claim 1 comprising
 - (a) 0.01 to 5% (w/v) sabeluzole polymorph I;
 - (b) 0.2 to 4% (w/v) suspending agents;
 - (c) up to 30% (w/v) sweeteners;
 - (d) up to 30% (v/v) preservatives;
 - (e) 0.01 to 1% (w/v) wetting agent;
 - (f) buffer substances to fix the pH in the range from 8 to 10; and
 - (g) water q.s. ad 100%.
10. A suspension according to claim 9 comprising:
 - (a) 0.05 to 1% (w/v) sabeluzole polymorph I;
 - (b) 0.8 to 1.5% (w/v) dispersible cellulose and optionally 0.1 to 0.5% (w/v) hypromellose;
 - (c) 10 to 30% (v/v) sorbitol solution;
 - (d) 10 to 30% (v/v) propylene glycol;
 - (e) 0.01 to 1% (w/v) polysorbate 20;

-12-

- (f) buffer substances to fix the pH in the range from 8.5 to 9.5; and
- (g) water q.s. ad 100%.

- 11. A suspension according to claim 10 comprising
 - 5 (a) 0.1% (w/v) sabeluzole polymorph I;
 - (b) 1.2% (w/v) dispersible cellulose and optionally 0.25% (w/v) hypromellose;
 - (c) 20% (v/v) sorbitol 70% (w/w) solution;
 - (d) 20% (v/v) propylene glycol;
 - (e) 0.025% (w/v) polysorbate 20;
 - 10 (f) sodium carbonate and hydrochloric acid to fix the pH at about 9;
 - (g) water q.s. ad 100%.
- 12. A process for preparing a suspension as claimed in any of claims 1 to 11
 - 15 characterized in that a therapeutically effective amount of sabeluzole is intimately mixed with a pharmaceutically acceptable carrier.
- 13. A process according to claim 12 comprising the following steps:
 - (a) the suspending agents, wetting agents, preservatives and sweeteners are mixed with an amount of water;
 - 20 (b) sabeluzole is mixed with phase (a); and
 - (c) the pH is fixed in the range from 8 to 10.
- 14. A suspension according to any of claims 1 to 11 for use as a medicine.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 95/03966

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/445 A61K9/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A,4 861 785 (R.A.STOKBROEKS ET AL.) 29 August 1989 cited in the application see claims 1,7-18 see examples 47,48 ---	1-14
A	EP,A,0 501 552 (JANSSEN PHARMACEUTICA N.V.,BE) 2 September 1992 see claims see examples 11-13 ---	1-14
A,P	WO,A,94 25029 (JANSSEN PHARMACEUTICA N.V.,BE) 10 November 1994 see claims 7,8,10 see example 2 -----	1-14

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

1 February 1996

Date of mailing of the international search report

16.02.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Scarponi, U

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 95/03966

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4861785	29-08-89	US-A- 5010198	23-04-91
		AU-B- 582601	06-04-89
		AU-B- 5057485	17-07-86
		BG-B- 60431	31-03-95
		CA-A- 1260474	26-09-89
		CY-A- 1690	14-01-94
		EP-A, B 0184257	11-06-86
		IE-B- 58807	17-11-93
		JP-B- 7059578	28-06-95
		JP-A- 61137884	25-06-86
		SU-A- 1428203	30-09-88
EP-A-501552	02-09-92	AT-T- 131478	15-12-95
		AU-B- 649788	02-06-94
		AU-B- 1208592	15-09-92
		BG-A- 98068	30-06-94
		CN-A- 1064865	30-09-92
		DE-D- 69206787	25-01-96
		WO-A- 9214731	03-09-92
		EP-A- 0573473	15-12-93
		HU-A- 67695	28-04-95
		JP-T- 6505012	09-06-94
		NZ-A- 241592	26-07-94
		NZ-A- 248437	26-07-94
		US-A- 5434168	18-07-95
		ZA-A- 9201341	24-08-93
WO-A-9425029	10-11-94	AU-B- 6678594	21-11-94
		CA-A- 2160365	10-11-94